

# 2021 Western Drought & Extreme Heat Assessment

Assessment Period: Aug 2-8, 2021

Publication Date: August 12, 2021

USDA NASS

Disaster Monitoring Team



# Outline

- The attached slides provide an overview of the extreme heat and drought conditions in five NASS Regions: Northwest, Pacific, Mountain, Northern Plains, and Upper Midwest.
  - Slides 3-5 illustrate **temperature and precipitation anomalies** for the conterminous U.S. from August 1-11, 2021. This is based on PRISM Climate Group data and 30 years of climatological information.
  - Slides 6-11 illustrate areas impacted by **heat stress** for each region individually for Weeks 30 (Jul 26-Aug 1, 2021) & 31 (Aug 2-8, 2021) in 2021, Week 31 in 2020, and the Week 31 five-year average.
  - Slides 12-30 identify the resulting impact of the lack of precipitation and extreme heat on **cropland subsoil moisture**. Weekly average subsoil moisture, anomalies, and categorical levels for Week 31 (Aug 2-8, 2021) are illustrated. The information was obtained from the Crop-CASMA web application. Figures use a crop mask (gray) to block out non-cropland areas. An analysis was conducted to identify the percent of cropland at varying levels with extreme conditions highlighted.

# PRISM Climate Group Data

- Offers an "early glimpse" version of precipitation and temperature data from the current month
- The datasets are modeled using climatologically-aided interpolation (CAI), which uses the long-term average pattern (i.e., the 30-year normals) as first-guess of the spatial pattern of climatic conditions for a given month or day
- Data supported by USDA RMA



Map provided by PRISM Climate Group: <https://prism.oregonstate.edu/mtd/>




# Total Precipitation Anomaly: 01 Aug 2021 - 11 Aug 2021

Period ending 7 AM EST 11 Aug 2021

Base period: 1981-2010

(Map created 12 Aug 2021)



Copyright (c) 2021, PRISM Climate Group, Oregon State University



Map provided by PRISM Climate Group: <https://prism.oregonstate.edu/mtd/>




# Daily Mean Temperature Anomaly: 01 Aug 2021 - 11 Aug 2021

Period ending 7 AM EST 11 Aug 2021

Base period: 1981-2010

(Map created 12 Aug 2021)



Copyright (c) 2021, PRISM Climate Group, Oregon State University



Map provided by PRISM Climate Group: <https://prism.oregonstate.edu/mtd/>



# Heat Stress Data

- Data calculated using data from two main sources of gridded products, PRISM, and RTMA.
- Heat stress is calculated as the difference between the maximum observed temperature during the day and the selected threshold ( $T_{dth}$ ). If the maximum temperature is lower than  $T_{dth}$ , HSDD is equal to zero.


$$HSDD = (T_{max} - T_{dth})$$



Source: NASS Climate-based Information System




### Northwest Region - Heat Index (93°F) - 2021: Week 31 Accumulated Degrees above 93 Degrees



\*Does not include Alaska

°F 32.3 33.3 34.6 36.9 41.7 48.2 57.9 80.6 113

### Northwest Region - Heat Index (97°F) - 2021: Week 31 Accumulated Degrees above 97 Degrees



\*Does not include Alaska


°F 32.3 33.3 34.6 36.9 41.7 48.2 57.9 80.6




Source: NASS Climate-based Information System



Pacific Region - Heat Index (93°F) - 2021: Week 31  
Accumulated Degrees above 93 Degrees




Pacific Region - Heat Index (97°F) - 2021: Week 31  
Accumulated Degrees above 97 Degrees



\*Does not include Hawaii



\*Does not include Hawaii




Source: NASS Climate-based Information System




### Mountain Region - Heat Index (93°F) - 2021: Week 31 Accumulated Degrees above 93 Degrees

2021: Week 31      2021: Week 30




2020: Week 31      Last 5 Years Average: Week 31





### Mountain Region - Heat Index (97°F) - 2021: Week 31 Accumulated Degrees above 97 Degrees

2021: Week 31      2021: Week 30




2020: Week 31      Last 5 Years Average: Week 31




Source: NASS Climate-based Information System



### Northern Plains Region - Heat Index (90°F) - 2021: Week 31 Accumulated Degrees above 90 Degrees






### Northern Plains Region - Heat Index (93°F) - 2021: Week 31 Accumulated Degrees above 93 Degrees






Source: NASS Climate-based Information System



### Upper Midwest Region - Heat Index (90°F) - 2021: Week 31 Accumulated Degrees above 90 Degrees



### Upper Midwest Region - Heat Index (93°F) - 2021: Week 31 Accumulated Degrees above 93 Degrees



Source: NASS Climate-based Information System



# Soil Moisture Data

- Hosted by Crop-CASMA (Crop Condition and Soil Moisture Analytics) <https://nassgeo.csiss.gmu.edu/CropCASMA/>
- Data Used
  - Sub Soil Moisture, 9km, Weekly, Year 2021, Week 31, Aug 2-8, 2021
  - Sub Soil Moisture Anomaly, 9km, Weekly, Year 2021, Week 31, Aug 2-8, 2021
  - Sub Soil Moisture Categorical, 9km, Weekly, Year 2021, Week 31, Aug 2-8, 2021
- Total Cropland derived by 2020 Cultivated Layer hosted on Crop-CASMA.



# Sub Soil Moisture

- NASA Remotely Sensed Rootzone Soil (sub soil) is defined as the top 3.2 feet (approximately 1 meter).
- The NASA SMAP (Soil Moisture Active Passive) 9km soil moisture measurements are volumetric soil moisture (i.e. volumetric water content in the soil). It is simply the ratio of water volume to soil volume.
- Sub soil moisture measuring at  $0.1 \text{ cm}^3/\text{cm}^3$  and below (10% water content) could be considered very dry.



# Sub Soil Moisture Anomaly

- The soil moisture anomaly (SMA) in CropCASMA is a measure of deviation of the current soil moisture value from the "normal" soil moisture level, which is represented by a historical average soil moisture value (from 2015 to current).
- The SMA of a given location is defined by the following formula:

$$SMA = \frac{SM - SM_m}{SM_m} \times 100\%$$

where SM and SM<sub>m</sub> denote current soil moisture value and the historical average soil moisture value of a given location.

- Soil moisture anomaly below -40% could be considered very abnormal, which means there is 40% less soil moisture than normal conditions.

# Sub Soil Moisture Categorical

- SMAP values are categorized into NASS categories which include:
  - Very Short - Soil moisture supplies are significantly less than what is required for normal plant development. Growth has been stopped or nearly so and plants are showing visible signs of moisture stress. Under these conditions, plants will quickly suffer irreparable damage.
  - Short - Soil dry. Seed germination and/or normal crop growth and development would be curtailed.
  - Adequate - Soil moist. Seed germination and/or crop growth and development would be normal or unhindered.
  - Surplus - Soil wet. Fields may be muddy and will generally be unable to absorb additional moisture. Young developing crops may be yellowing from excess moisture.




Northwest Region  
 Sub Soil Moisture 9km  
 Aug 2-8, 2021



NW Sub SM 9km Aug 2-8, 2021



Sub Soil Moisture (9km, Aug 2-8, 2021)				
Volumetric Soil Moisture (cm <sup>3</sup> /cm <sup>3</sup> )	Northwest Region	Idaho	Oregon	Washington
	Percentage of Total Cropland			
0.0-0.05	4.69%	2.90%	5.71%	5.74%
0.05-0.1	17.60%	18.18%	6.55%	22.25%
0.1-0.15	25.57%	27.82%	18.18%	27.52%
0.15-0.2	36.81%	34.50%	36.29%	38.46%
0.2-0.25	11.62%	15.81%	18.71%	4.93%
0.25-0.3	2.63%	0.80%	10.73%	0.40%
0.3-0.35	0.79%	0.00%	3.22%	0.33%
0.35-0.4	0.10%	0.00%	0.50%	0.00%
0.4-0.45	0.02%	0.00%	0.11%	0.00%
0.45-0.5	0.00%	0.00%	0.00%	0.00%
0.5-0.55	0.00%	0.00%	0.00%	0.00%
0.55-0.6	0.00%	0.00%	0.00%	0.00%
0.6-0.65	0.16%	0.00%	0.00%	0.37%
> 0.65	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScope - <http://nassgeodata.gmu.edu/VegScope>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Northwest Region  
 Sub Soil Moisture Anomaly 9km  
 Aug 2-8, 2021



NW Sub SM Anomaly 9km Aug 2-8, 2021



Sub Soil Moisture Anomaly (9km, Aug 2-8, 2021)				
Soil Moisture Anomaly	Northwest Region	Idaho	Oregon	Washington
	Percentage of Total Cropland			
<-70%	0.00%	0.00%	0.00%	0.00%
-70%~-50%	0.00%	0.00%	0.00%	0.00%
-50%~-40%	0.00%	0.00%	0.00%	0.00%
-40%~-30%	3.55%	5.36%	1.83%	2.91%
-30%~-20%	12.41%	6.03%	8.27%	19.79%
-20%~-10%	51.78%	34.47%	59.06%	62.73%
-10%~0%	26.98%	41.17%	28.09%	14.56%
0%~-10%	4.69%	11.33%	2.75%	0.01%
10%~20%	0.58%	1.62%	0.00%	0.00%
20%~30%	0.00%	0.01%	0.00%	0.00%
30%~40%	0.00%	0.00%	0.00%	0.00%
40%~50%	0.00%	0.00%	0.00%	0.00%
50%~70%	0.00%	0.00%	0.00%	0.00%
>70%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Northwest Region  
 Sub Soil Moisture Categorical 9km  
 Aug 2-8, 2021

Sub Soil Moisture Categorical (9km, Aug 2-8, 2021)				
Categorical Soil Moisture	Northwest Region	Idaho	Oregon	Washington
	Percentage of Total Cropland			
Very Short	49.68%	34.87%	59.76%	57.48%
Short	30.69%	27.48%	30.54%	33.53%
Adequate	18.90%	37.65%	8.84%	7.70%
Surplus	0.53%	0.00%	0.86%	0.82%
No Data	0.21%	0.00%	0.00%	0.48%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



NW Sub SM Category 9km Aug 2-8, 2021



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Pacific Region  
 Sub Soil Moisture 9km  
 Aug 2-8, 2021

Sub Soil Moisture (9km, Aug 2-8, 2021)			
Volumetric Soil Moisture	Pacific Region	California	Nevada
	Percentage of Total Cropland	Percentage of Total Cropland	Percentage of Total Cropland
0.0-0.05	8.42%	8.05%	17.31%
0.05-0.1	21.09%	21.19%	17.91%
0.1-0.15	19.63%	18.98%	35.13%
0.15-0.2	20.40%	20.32%	22.04%
0.2-0.25	29.20%	30.18%	6.96%
0.25-0.3	0.85%	0.86%	0.65%
0.3-0.35	0.40%	0.42%	0.00%
0.35-0.4	0.00%	0.00%	0.00%
0.4-0.45	0.00%	0.00%	0.00%
0.45-0.5	0.00%	0.00%	0.00%
0.5-0.55	0.00%	0.00%	0.00%
0.55-0.6	0.00%	0.00%	0.00%
0.6-0.65	0.00%	0.00%	0.00%
> 0.65	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Pacific Sub SM 9km Aug 2-8, 2021



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Pacific Region  
 Sub Soil Moisture Anomaly 9km  
 Aug 2-8, 2021



Pacific Sub SM Anomaly 9km Aug 2-8, 2021



Sub Soil Moisture Anomaly (9km, Aug 2-8, 2021)			
Soil	Pacific Region	California	Nevada
Moisture Anomaly	Percentage of Total Cropland	Percentage of Total Cropland	Percentage of Total Cropland
<-70%	0.00%	0.00%	0.00%
-70%~-50%	0.63%	0.06%	13.67%
-50%~-40%	1.80%	1.64%	5.50%
-40%~-30%	1.53%	1.29%	6.66%
-30%~-20%	3.47%	2.95%	15.16%
-20%~-10%	27.96%	27.61%	36.27%
-10%~0%	61.69%	63.50%	20.20%
0%~-10%	2.92%	2.94%	2.47%
10%~20%	0.00%	0.00%	0.10%
20%~30%	0.00%	0.00%	0.00%
30%~40%	0.00%	0.00%	0.00%
40%~50%	0.00%	0.00%	0.00%
50%~70%	0.00%	0.00%	0.00%
>70%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScope - <http://nassgeodata.gmu.edu/VegScope>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Pacific Region  
 Sub Soil Moisture Categorical 9km  
 Aug 2-8, 2021

Sub Soil Moisture Categorical (9km, Aug 2-8, 2021)			
Categorical Soil Moisture	Pacific Region Percentage of Total Cropland	California Percentage of Total Cropland	Nevada Percentage of Total Cropland
Very Short	56.32%	56.92%	41.90%
Short	19.59%	19.28%	27.18%
Adequate	23.43%	23.33%	25.65%
Surplus	0.21%	0.01%	5.27%
No Data	0.45%	0.47%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Pacific Sub SM Category 9km Aug 2-8, 2021



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>




Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>



# Mountain Region Sub Soil Moisture 9km Aug 2-8, 2021

Sub Soil Moisture (9km, Aug 2-8, 2021)							
Volumetric Soil Moisture (cm <sup>3</sup> /cm <sup>3</sup> )	Mountain Region	Arizona	Colorado	Montana	New Mexico	Utah	Wyoming
	Percentage of Total Cropland						
0.0-0.05	1.73%	4.52%	3.01%	0.28%	0.04%	2.67%	7.39%
0.05-0.1	6.20%	32.34%	7.06%	1.17%	10.30%	6.96%	19.05%
0.1-0.15	12.11%	14.03%	16.16%	6.84%	8.99%	23.65%	33.46%
0.15-0.2	38.18%	38.31%	32.57%	41.52%	41.53%	43.75%	25.99%
0.2-0.25	32.45%	7.24%	39.07%	34.93%	27.49%	21.32%	10.79%
0.25-0.3	9.12%	3.40%	2.12%	15.04%	10.43%	1.66%	2.97%
0.3-0.35	0.21%	0.17%	0.01%	0.23%	1.21%	0.00%	0.35%
0.35-0.4	0.00%	0.00%	0.00%	0.00%	0.01%	0.00%	0.00%
0.4-0.45	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
0.45-0.5	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
0.5-0.55	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
0.55-0.6	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
0.6-0.65	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
> 0.65	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>						




Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>



# Mountain Region Sub Soil Moisture Anomaly 9km Aug 2-8, 2021

Sub Soil Moisture Anomaly (9km, Aug 2-8, 2021)							
Soil Moisture Anomaly	Mountain Region	Arizona	Colorado	Montana	New Mexico	Utah	Wyoming
	Percentage of Total Cropland						
<-70%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
-70%~-50%	0.08%	0.00%	0.00%	0.08%	0.00%	0.37%	0.48%
-50%~-40%	0.35%	1.12%	0.30%	0.11%	0.00%	0.12%	3.24%
-40%~-30%	1.16%	2.28%	1.33%	0.42%	0.06%	0.68%	9.13%
-30%~-20%	6.62%	2.06%	7.79%	6.51%	2.09%	2.57%	17.15%
-20%~-10%	43.14%	12.41%	39.39%	52.87%	25.01%	22.23%	44.01%
-10%~0%	39.82%	27.74%	44.21%	37.21%	44.97%	60.74%	24.37%
0%~-10%	6.24%	20.84%	6.45%	2.79%	18.66%	12.46%	1.62%
10%~20%	1.43%	12.71%	0.51%	0.00%	9.01%	0.84%	0.00%
20%~30%	1.10%	19.84%	0.03%	0.00%	0.21%	0.00%	0.00%
30%~40%	0.05%	0.99%	0.00%	0.00%	0.00%	0.00%	0.00%
40%~50%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
50%~70%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
>70%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>						




Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>



Mountain Region  
 Sub Soil Moisture Categorical 9km  
 Aug 2-8, 2021

Sub Soil Moisture Categorical (9km, Aug 2-8, 2021)							
Categorical Soil Moisture	Mountain Region	Arizona	Colorado	Montana	New Mexico	Utah	Wyoming
	Percentage of Total Cropland						
Very Short	39.42%	0.00%	13.28%	61.85%	28.26%	28.17%	34.28%
Short	32.21%	0.00%	33.85%	32.34%	41.19%	30.25%	48.11%
Adequate	25.50%	66.44%	52.87%	3.96%	30.31%	40.81%	15.88%
Surplus	2.07%	33.54%	0.00%	0.25%	0.23%	0.70%	1.74%
No Data	0.80%	0.02%	0.00%	1.59%	0.01%	0.06%	0.00%
<b>Total</b>	<b>100.00%</b>						



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Northern Plains Region  
Sub Soil Moisture 9km  
Aug 2-8, 2021



N.Plains Sub SM 9km Aug 2-8, 2021



Sub Soil Moisture (9km, Aug 2-8, 2021)					
Volumetric Soil Moisture (cm3/cm3)	Northern Plains Region	Kansas	Nebraska	North Dakota	South Dakota
	Percentage of Total Cropland				
0.0-0.05	0.64%	0.02%	2.54%	0.16%	0.03%
0.05-0.1	4.17%	4.09%	10.89%	1.47%	0.42%
0.1-0.15	10.89%	11.53%	11.85%	13.25%	5.65%
0.15-0.2	35.49%	22.86%	23.38%	62.89%	29.74%
0.2-0.25	27.72%	30.69%	23.20%	16.64%	43.81%
0.25-0.3	18.08%	22.43%	27.12%	4.49%	20.17%
0.3-0.35	2.65%	7.28%	0.89%	1.11%	0.18%
0.35-0.4	0.35%	1.11%	0.12%	0.00%	0.00%
0.4-0.45	0.00%	0.00%	0.00%	0.00%	0.00%
0.45-0.5	0.00%	0.00%	0.00%	0.00%	0.00%
0.5-0.55	0.00%	0.00%	0.00%	0.00%	0.00%
0.55-0.6	0.00%	0.00%	0.00%	0.00%	0.00%
0.6-0.65	0.00%	0.00%	0.00%	0.00%	0.00%
> 0.65	0.00%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Northern Plains Region  
 Sub Soil Moisture Anomaly 9km  
 Aug 2-8, 2021



N.Plains Sub SM Anomaly 9km Aug 2-8, 2021



Sub Soil Moisture Anomaly (9km, Aug 2-8, 2021)					
Soil Moisture Anomaly	Northern Plains Region	Kansas	Nebraska	North Dakota	South Dakota
	Percentage of Total Cropland				
<-70%	0.00%	0.00%	0.00%	0.00%	0.00%
-70%~-50%	0.02%	0.00%	0.09%	0.00%	0.00%
-50%~-40%	0.56%	0.32%	1.84%	0.11%	0.07%
-40%~-30%	1.83%	1.65%	2.77%	2.45%	0.18%
-30%~-20%	24.58%	16.84%	14.14%	43.64%	21.68%
-20%~-10%	54.16%	64.10%	50.06%	38.71%	65.49%
-10%~0%	16.74%	12.06%	28.27%	15.09%	12.58%
0%~-10%	2.11%	5.04%	2.83%	0.00%	0.00%
10%~20%	0.00%	0.00%	0.00%	0.00%	0.00%
20%~30%	0.00%	0.00%	0.00%	0.00%	0.00%
30%~40%	0.00%	0.00%	0.00%	0.00%	0.00%
40%~50%	0.00%	0.00%	0.00%	0.00%	0.00%
50%~70%	0.00%	0.00%	0.00%	0.00%	0.00%
>70%	0.00%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>



Northern Plains Region  
 Sub Soil Moisture Categorical 9km  
 Aug 2-8, 2021




N.Plains Sub SM Category 9km Aug 2-8, 2021



SMAP-CATEGORY Legend

- Very Short
- Short
- Adequate
- Surplus
- No Data



Produced by VegScope - <http://nassgeodata.gmu.edu/VegScope>

Sub Soil Moisture Categorical (9km, Aug 2-8, 2021)					
Categorical Soil Moisture	Northern Plains Region	Kansas	Nebraska	North Dakota	South Dakota
	Percentage of Total Cropland				
Very Short	23.93%	5.02%	10.09%	42.85%	40.71%
Short	39.92%	32.38%	46.46%	41.38%	41.20%
Adequate	35.09%	59.87%	43.45%	14.78%	18.05%
Surplus	0.79%	2.73%	0.00%	0.00%	0.03%
No Data	0.28%	0.00%	0.00%	0.99%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Upper Midwest Region  
Sub Soil Moisture 9km  
Aug 2-8, 2021

Sub Soil Moisture (9km, Aug 2-8, 2021)				
Volumetric Soil Moisture (cm3/cm3)	Upper Midwest Region	Iowa	Minnesota	Wisconsin
	Percentage of Total Cropland			
0.0-0.05	0.38%	0.00%	0.96%	0.08%
0.05-0.1	4.07%	0.05%	6.17%	10.08%
0.1-0.15	7.23%	0.00%	12.53%	14.59%
0.15-0.2	13.40%	3.21%	24.39%	16.20%
0.2-0.25	37.51%	35.77%	44.18%	26.28%
0.25-0.3	28.54%	46.73%	9.25%	23.71%
0.3-0.35	7.80%	13.07%	2.20%	6.54%
0.35-0.4	0.75%	1.18%	0.22%	0.81%
0.4-0.45	0.07%	0.00%	0.09%	0.20%
0.45-0.5	0.11%	0.00%	0.00%	0.64%
0.5-0.55	0.05%	0.00%	0.00%	0.28%
0.55-0.6	0.04%	0.00%	0.00%	0.26%
0.6-0.65	0.03%	0.00%	0.00%	0.19%
> 0.65	0.02%	0.00%	0.00%	0.14%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



U.Midwest Sub SM 9km Aug 2-8, 2021



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>



Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>




Upper Midwest Region  
 Sub Soil Moisture Anomaly 9km  
 Aug 2-8, 2021

Sub Soil Moisture Anomaly (9km, Aug 2-8, 2021)				
Soil Moisture Anomaly	Upper Midwest Region	Iowa	Minnesota	Wisconsin
	Percentage of Total Cropland			
<-70%	0.00%	0.00%	0.00%	0.00%
-70%~-50%	0.16%	0.00%	0.42%	0.00%
-50%~-40%	0.90%	0.00%	2.35%	0.07%
-40%~-30%	2.89%	0.00%	6.73%	2.00%
-30%~-20%	33.41%	13.64%	63.62%	18.37%
-20%~-10%	34.13%	40.47%	26.14%	34.66%
-10%~0%	22.28%	36.72%	0.73%	32.50%
0%~-10%	5.87%	8.52%	0.00%	12.03%
10%~20%	0.29%	0.66%	0.00%	0.00%
20%~30%	0.06%	0.00%	0.00%	0.38%
30%~40%	0.00%	0.00%	0.00%	0.00%
40%~50%	0.00%	0.00%	0.00%	0.00%
50%~70%	0.00%	0.00%	0.00%	0.00%
>70%	0.00%	0.00%	0.00%	0.00%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



U.Midwest Sub SM Anomaly 9km Aug 2-8, 2021



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>




Crop-CASMA: <https://nassgeo.csiss.gmu.edu/CropCASMA/>



Upper Midwest Region  
Sub Soil Moisture Categorical 9km  
Aug 2-8, 2021

Sub Soil Moisture Categorical (9km, Aug 2-8, 2021)				
Categorical Soil Moisture	Upper Midwest Region	Iowa	Minnesota	Wisconsin
	Percentage of Total Cropland			
Very Short	22.13%	16.40%	34.20%	10.46%
Short	39.09%	43.10%	43.60%	18.88%
Adequate	37.10%	40.51%	22.14%	61.05%
Surplus	1.60%	0.00%	0.00%	9.31%
No Data	0.08%	0.00%	0.07%	0.31%
<b>Total</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>	<b>100.00%</b>



Produced by VegScape - <http://nassgeodata.gmu.edu/VegScape>